



# ENVIRONMENTAL HEALTH PERSPECTIVES

<http://www.ehponline.org>

## Acute Impact of Hourly Ambient Air Pollution on Preterm Birth

Shanshan Li, Yuming Guo, and Gail Williams

<http://dx.doi.org/10.1289/EHP200>

**Received: 23 June 2015**

**Revised: 29 October 2015**

**Accepted: 15 April 2016**

**Published: 29 April 2016**

**Note to readers with disabilities:** *EHP* will provide a [508-conformant](#) version of this article upon final publication. If you require a 508-conformant version before then, please contact [ehp508@niehs.nih.gov](mailto:ehp508@niehs.nih.gov). Our staff will work with you to assess and meet your accessibility needs within 3 working days.



National Institute of  
Environmental Health Sciences

# **Acute Impact of Hourly Ambient Air Pollution on Preterm Birth**

Shanshan Li, Yuming Guo, and Gail Williams

School of Public Health, The University of Queensland, Brisbane, Queensland, Australia

**Address correspondence to** Shanshan Li, School of Public Health, The University of Queensland, Herston Road, Herston, Brisbane, QLD 4006, Australia. Telephone: +61 7 3346 4639. E-mail: [uqshandy0601@gmail.com](mailto:uqshandy0601@gmail.com)

**Running title:** Hourly air pollution and preterm birth

**Acknowledgments:** S. L. is supported by the Centre for Air quality & health Research and evaluation, Australia. Y. G. is supported by The University of Queensland Postdoctoral Research Fellowship.

**Conflict of Interest:** None declared.

## **Abstract**

**Background:** Preterm birth is a major perinatal health problem but factors leading to it are still not completely understood.

**Objectives:** To identify the relation between acute increase in ambient air pollution in a few hours before onset of labour and the risk of preterm birth.

**Methods:** We collected registered birth outcome data and hourly ambient air pollution measurements during 2009 – 2013 in Brisbane, Australia. Using the time-stratified case-crossover design and conditional logistic regression models with natural cubic splines, we assessed the shape of air pollution – preterm birth curve, after controlling for potential confounders. We also examined the effect modification of other factors.

**Results:** The association between air pollution [nitrogen dioxide (NO<sub>2</sub>), sulphur dioxide (SO<sub>2</sub>), and carbon monoxide (CO)] and preterm birth was non-linear. Threshold concentrations for the mean of 0 – 24 hours NO<sub>2</sub>, 24 – 48 hours SO<sub>2</sub>, and 24 – 48 hours CO before onset of labour were 7.6 parts per billion (ppb), 3.8 ppb, and 162.5 ppb, respectively. Increases in air pollution concentrations above thresholds were associated with increased risks of preterm birth. The ORs of preterm birth at 95th percentile of NO<sub>2</sub>, SO<sub>2</sub> and CO against the thresholds were 1.17 (1.08, 1.27), 1.01 (0.99, 1.04), and 1.18 (1.06, 1.32), respectively. The associations were modified by demographic factors, for example, maternal smoking and socioeconomic status (SES).

**Conclusion:** Acute increases in ambient air pollution concentrations above certain levels before onset of labour may stimulate preterm birth.

## **Introduction**

Preterm birth is a major perinatal health problem associated with neonatal mortality and morbidity and can cause long-term adverse health consequences in life (Beck et al. 2010; Lumley 2003). It was estimated that around 11% of all live births were born preterm worldwide in 2010, and this high and still rising incidence represents significant financial implications for healthcare systems (Blencowe et al. 2012). Yet, factors leading to preterm birth are still not completely understood. In recent years, there has been a growing concern about the possible influence of air pollution on preterm birth. The majority of studies (Bobak 2000; Gehring et al. 2011; C Hansen et al. 2006; Huynh et al. 2006; Jalaludin et al. 2007; Leem et al. 2006; Llop et al. 2010; Ritz et al. 2000; Ritz et al. 2007; Suh et al. 2008; Suh et al. 2009; Wu et al. 2009) investigated this relationship focused on exposures to air pollutants during the entire pregnancy or during specific trimesters, and the results were mixed. We have found no published studies on the impact of maternal exposure to air pollution in the few hours before onset of labour on the risk of preterm birth.

The aim of the present study was to identify the relation between maternal exposures to ambient air pollutants a few hours before onset of labour and preterm birth in Brisbane, Australia, from 2009 to 2013. We hypothesized that a short-term increase in ambient air pollution closely before onset of labour contributes to the risk of preterm birth and that such effect would be modified by demographic factors such as maternal age, maternal existing medical conditions, number of previous pregnancies, smoking status, multiple birth, baby gender, and socioeconomic levels.

## **Materials and Methods**

### ***Birth Information***

We collected birth outcome data from the Queensland Health Perinatal Data Collection Unit for births during 1 January 2009 – 31 December 2013 in Brisbane metropolitan area, Australia. It is a registered database covers births from all public and private hospitals and voluntarily reported homebirths in Brisbane. Information used in this study includes birth status (born alive or stillbirth), gestational age in weeks, date and time of birth, method of birth, length of labour, number of births, baby gender, maternal age, pre-pregnancy medical conditions, number of previous pregnancies, maternal smoking status, and an index of socioeconomic status (SES) linked to the living area during pregnancy. The definition of preterm birth is birth before a gestational age of 37 weeks (Goldenberg et al. 2008). Thus, we retained data on all live births occurring less than 37 weeks of gestation in this study. The study was approved by the University of Queensland Medical Research Ethics Committee.

### ***Air Pollution and Meteorological Data***

Hourly data on particulate matter 10 micrometres or less in diameter (PM<sub>10</sub>), particulate matter 2.5 micrometres or less in diameter (PM<sub>2.5</sub>), nitrogen dioxide (NO<sub>2</sub>), sulphur dioxide (SO<sub>2</sub>), ozone (O<sub>3</sub>), carbon monoxide (CO), ambient temperature, and relative humidity were obtained from the Queensland Government Department of Environment and Heritage Protection for 5 monitoring sites across Brisbane. PM<sub>2.5</sub> and PM<sub>10</sub> were measured by either a high or low volume air sampler or a tapered element oscillating microbalance (TEOM). NO<sub>2</sub> was measured with a technique known as “chemiluminescence”, which is a chemical reaction that emits energy in the form of light. SO<sub>2</sub> was measured by a Differential Optical Absorbance Spectroscopy (DOAS) instrument. O<sub>3</sub> concentrations were monitored through the principle of absorption of ultraviolet (UV) light. CO was measured by a technique known as “gas filter correlation”.

For each preterm birth, we defined the onset of labour time as the birth time by a deduction of labour length. Then we calculated mean values of air pollutants and meteorological conditions by hourly measurements for the periods of 0 – 24 hours, 24 – 48 hours, 48 – 72 hours, 0 – 48 hours, and 0 – 72 hours, respectively, before the time of labour onset if at least 75% of measurements were available in the corresponding period. Otherwise the data was considered missing data.

### ***Statistical Analysis***

We employed a time-stratified case-crossover design to examine the association between air pollution and preterm birth. The case-crossover design can be explained as a self-matched case-control study (Janes et al. 2005). For each individual, exposure information (e.g., air pollution) is collected for the “case” period (that is, the onset of labour time of preterm birth in this study) and a series of “control” periods that are not associated with the event of interest. In the time-stratified design, control periods should be selected from the fixed time strata (e.g., month) to avoid any “overlap bias” (Lumley and Levy 2000). In this study, we used the calendar month as the fixed time strata, and control periods comprised the same hour of the same day of the week in the calendar month of preterm birth labour onset, to control for the effect of day of the week and intra-day variation. Air pollution and confounding information were obtained for the hour of the onset of labour event for both case and control periods.

We used conditional logistic regression models to fit the time-stratified case-crossover design, which successfully controlled for time-invariant individual level confounders (e.g., baby gender and maternal age), because comparisons between case and control periods were made within individuals. Mean values of each air pollutant (PM<sub>10</sub>, PM<sub>2.5</sub>, NO<sub>2</sub>, SO<sub>2</sub>, O<sub>3</sub>,

and CO) for the periods of 0 – 24 hours, 24 – 48 hours, 48 – 72 hours, 0 – 48 hours, and 0 – 72 hours, respectively, preceding the time of labour onset were examined separately in single-pollutant models. To fully adjust for the potential time-variant confounders, we used natural cubic splines with 4 degrees of freedom (df) for the 0 – 72 hours mean values of ambient temperature and relative humidity in all models.

Natural cubic splines were also applied for air pollutants in single-pollutant models to check whether the associations between air pollutants and preterm birth were linear or non-linear. We selected the df and the time frame of exposure prior to onset of labour by judging the model fit which is reflected by the Akaike Information Criterion (AIC). For each air pollutant, model with the lowest AIC value indicated the best df and exposure time frame. In case of linear relationship between air pollution and preterm birth, we would calculate the odds ratios (OR) and the 95% confidence intervals (CI) of preterm birth at 75th and 95th percentiles of air pollution against the median concentration of air pollution. Otherwise in case of non-linear relationship, we would calculate the OR and 95% CI of preterm birth at 75th and 95th percentiles of air pollution against the minimum-preterm birth concentration of air pollution (threshold).

In order to determine the threshold of air pollutant, we first plotted the graph of the relationship between air pollutant and preterm birth, and then visually checked the possible range of the threshold. Afterwards, we iteratively estimated the AIC values for conditional logistic regression models by 0.1 unit increment in air pollutant within the identified range of threshold from visual inspection using the segment spline model. The concentrations of air pollutants corresponding to the lowest AIC values were chosen as the thresholds (minimum-preterm birth concentrations of air pollutants). This method has been widely used to test the

threshold for non-linear temperature effect on mortality (Chung et al. 2009; Kim et al. 2006; Yu et al. 2010).

To evaluate the potential confounding effects of other air pollutants on the association between an air pollutant and preterm birth, we ran two-pollutant and multiple-pollutant models, and compared the OR and 95% CI of preterm birth associated with a pollutant from the single-pollutant model with that from the two-pollutant or multiple-pollutant model. To assess the possible modification effects of demographic factors on the air pollution – preterm birth association, we conducted stratification analyses for different groups: maternal age (< 35 years VS  $\geq 35$  years), pre-pregnancy medical conditions (yes VS no), number of previous pregnancies (0 VS  $\geq 1$ ), maternal smoking status [smokers (self-reported any smoking during pregnancy) VS non-smokers (self-reported no smoking during pregnancy)], number of births (single birth VS multiple births), baby gender (girls VS boys), and SES index [1 – 5 (indicating low development) VS 6 – 10 (indicating high development)]. The statistical significance of difference between effect estimates for above subgroups (e.g., maternal age < 35 years VS  $\geq 35$  years), was examined by  $(\hat{Q}_1 - \hat{Q}_2) \pm 1.96\sqrt{S\hat{E}_1 + S\hat{E}_2}$ , where  $\hat{Q}_1$  and  $\hat{Q}_2$  are the effect estimates for the two categories (e.g., < 35 years and  $\geq 35$  years), and  $S\hat{E}_1$  and  $S\hat{E}_2$  are their respective standard errors (Kan et al. 2008; Zeka et al. 2006). Planned caesareans were excluded in the analyses.

We added temperature variability (standard deviation of 0–72 hours' temperatures) to the models, to check whether the effects of air pollutants on preterm birth would be changed or not, as studies reported that large temperature change increased risk of health events (Guo et al. 2011; Li et al. 2014; Qiu et al. 2013). We used 0–12 hours' average concentrations of air pollutants as exposure, to check whether the model fit was improved or not. All analyses



were performed using R software (version 3.1.3) (R Core Team 2015). R codes were provided to show how to match case and control by the same hour of the same day of the week in the same month (see Supplemental Material, R codes).

## **Results**

Table 1 shows the maternal and foetal demographic characteristics of preterm births delivered in Brisbane, Australia from January 1, 2009 to December 31, 2013. A total of 6949 preterm births occurred over the entire study period. Mothers younger than 35 years of age, without existing pre-pregnancy medical condition, ever got pregnant before the current pregnancy, never smoke, and mothers living in the area with high social-economic levels accounted for greater proportion of births. A majority of births were singletons and the baby genders distributed evenly.

Table 2 displays the hourly air pollution and meteorological exposure information during the study period. Mean levels of hourly PM<sub>2.5</sub>, PM<sub>10</sub>, NO<sub>2</sub>, SO<sub>2</sub>, O<sub>3</sub>, CO, ambient temperature, and relative humidity were 6.32 µg/m<sup>3</sup>, 17.27 µg/m<sup>3</sup>, 6.52 ppb, 1.95 ppb, 17.27 ppb, 219.30 ppb, 21.97 °C, and 70.82%, respectively. The summary statistics for daily air pollution and meteorological factors are similar as the hourly data (see Supplemental Material, Table S1). Hourly values of air pollutants and weather conditions were positively correlated with each other (see Supplemental Material, Table S2).

Our preliminary analyses (see Supplemental Material, Figure S1) showed that the associations between air pollutants and preterm birth were generally U-shaped for NO<sub>2</sub>, SO<sub>2</sub>, and CO, that is, the threshold effect (a minimum-preterm birth concentration of the air pollutant) was indicated. There were no statistically significant relationships among PM<sub>2.5</sub>,

PM10, O3 and preterm births in single-pollutant models. Natural cubic splines with 3df for the mean values of 0 – 24 hours NO2, 24 – 48 hours SO2, and 24 – 48 hours CO preceding the time of labour onset, respectively, produced the best model fits for each air pollutant. Thus, we chose these three air pollutants with certain exposure time frames to estimate the risk of preterm birth at 75th and 95th percentiles of pollutant against the threshold in all following analyses.

The threshold concentrations as listed in Table 3 were 7.6 ppb for the mean of antepartum 0 – 24 hours NO2, 3.8 ppb for the mean of antepartum 24 – 48 hours SO2, and 162.5 ppb for the mean of antepartum 24 – 48 hours CO, respectively. Figure 1 clearly presents the U-shaped relationships between air pollutants and preterm birth. Increased concentrations of NO2, SO2 and CO above thresholds shortly before onset of labour were positively associated with increased risks of preterm birth. OR values of preterm birth at 75th percentile of NO2, SO2 and CO against the thresholds were 1.01 (0.99, 1.03), 1.04 (0.99, 1.08), and 1.10 (1.01, 1.19), respectively; while OR values of preterm birth at 95th percentile of NO2, SO2 and CO against the thresholds were 1.17 (1.08, 1.27), 1.01 (0.99, 1.04), and 1.18 (1.06, 1.32), respectively (Table 3).

The effect of each pollutant on the risk of preterm birth appeared independent of the other pollutants (Figure 2). The OR values of preterm birth associated with individual air pollutants were similar in two-pollutant and three-pollutant models compared with those in single-pollutant models (Table 4).

Table 5 shows the results from stratification analyses for different groups evaluating whether demographic factors modified the air pollution – preterm birth relationship. There was no

consistent increasing or decreasing trend of the risk within categories of maternal age and pre-pregnancy medical conditions. The associations of NO<sub>2</sub> and CO with preterm birth were slightly stronger among those had ever got pregnant, multiple births, and female babies. The risks of preterm birth associated with all three air pollutants were greater among smokers and families living in lower SES areas. However, only smoking during pregnancy and low SES significantly modified the effects of NO<sub>2</sub> and CO on preterm birth, respectively (see Supplemental Material, Table S3).

The effects of air pollutants on preterm birth did not change when we put temperature variability to the models (see Supplemental Material, Figure S2). When we used 0–12 hours' average concentrations of air pollutants, the model fit was not improved.

## **Discussion**

In this study, we found the threshold effect (concentration of the air pollutant corresponding to minimum-preterm birth) of NO<sub>2</sub>, SO<sub>2</sub> and CO on preterm birth. Increased risks of preterm birth were associated with increases in the mean concentrations of NO<sub>2</sub>, SO<sub>2</sub> and CO above thresholds in 24 or 48 hours before birth. The effect of each pollutant appeared independent of the others. We did not discover effect modification with maternal age or maternal health history. The associations of air pollutants with preterm birth risk were stronger among smokers and women from areas with relatively lower SES, and the risks associated with NO<sub>2</sub> and CO were slightly greater among once pregnant women, multiple births, and female babies.

The shape of the exposure-response curve is a critical issue in air pollution researches. A threshold value of air pollutant effect is usually expected to protect population health by

keeping the pollutant below this level. Regarding the adverse effects of air pollution on birth outcomes, a recent analysis (Fleischer et al. 2014) of the World Health Organization Global Survey on Maternal and Perinatal Health suggested a possible threshold effect for preterm birth in China, with a threshold of 36.5 mg/m<sup>3</sup> for PM<sub>2.5</sub>. A Spanish study (Llop et al. 2010) observed that perinatal exposure to traffic-related air pollution above certain concentration levels was associated with preterm birth, with a threshold of 46.2 mg/m<sup>3</sup> for NO<sub>2</sub> and a threshold of 2.7 mg/m<sup>3</sup> for benzene. In our study, we again identified evidence for threshold effect of air pollution on preterm birth. However, our findings are related to short-term effects of air pollution on preterm birth rather than long-term effect.

Our findings of increased preterm birth associated with increases in NO<sub>2</sub>, SO<sub>2</sub> and CO concentrations in the immediate few hours preceding onset of labour are consistent with results of previous studies that reported the short-term effect of air pollution during pregnancy on preterm birth (Leem et al. 2006; Liu et al. 2003; Zhao et al. 2011). Liu et al (Liu et al. 2003) found that exposure to increased SO<sub>2</sub> and CO during the last month of pregnancy contributed to higher risk of preterm birth in Vancouver, Canada. Leem et al (Leem et al. 2006) detected dose dependent relationships between preterm birth and exposure to NO<sub>2</sub>, SO<sub>2</sub> and CO particularly during the third trimester of pregnancy in the Republic of Korea. Study (Zhao et al. 2011) conducted in Guangzhou, China also indicated positive associations between preterm birth and daily concentrations of NO<sub>2</sub> and SO<sub>2</sub> using a time-series design. Importantly, the concentrations of air pollutants in our study were lower than those of above studies. This means even people live in the environment with a very low concentration of air pollution, the air pollution still has impacts on human health. Some studies also found hazard effects of daily air pollution on daily preterm birth in US cities

which had slightly higher concentrations of air pollutants than our study area (Dar-Row et al. 2008; Darrow et al. 2009; Sagiv et al. 2005).

The very acute association between air pollution and the risk of preterm birth may suggest that ambient air pollutants can rapidly motivate the biologic mechanism of labour, leading to preterm birth in the next few hours. The potential mechanism responsible for this association may work through a series of causes including oxidative stress, inflammation, endothelial dysfunction, endocrine disruption, and hemodynamic responses (Cunningham et al. 2014; Slama et al. 2008). When pollutants are inhaled into the body, cytokines trigger oxidative stress which could induce endothelial dysfunction to develop preeclampsia (pregnancy hypertension) (Wu et al. 2009; Yorifuji et al. 2015). Simultaneously, air pollutants cause intrauterine inflammation which increases prostaglandin levels to progress preterm premature rupture of membranes (PPROM) (Aagaard-Tillery et al. 2005; Leem et al. 2006). Preterm birth is a consequence of preeclampsia and PPRM.

Interestingly, we didn't found significant impact of PM and O<sub>3</sub> on preterm birth in this study. A previous study conducted in the same study area reported that a long-term exposure to PM<sub>10</sub> and O<sub>3</sub> during trimester one was associated with an increased risk of preterm birth, with OR = 1.15 (95% CI: 1.06–1.25) and OR = 1.26 (95% CI: 1.10–1.45), respectively. This indicates that PM and O<sub>3</sub> might not have short-term effects on preterm birth, but have long-term (cumulative) effects (C. Hansen et al. 2006). Some studies using daily time series air pollution and preterm birth also found non-significant short-term impacts of PM (Darrow et al. 2009; Sagiv et al. 2005), but reported significant impacts of NO<sub>2</sub> and SO<sub>2</sub> (Olsson et al. 2012; Sagiv et al. 2005). Further studies are still needed to explore why PM and O<sub>3</sub> don't have short-term impacts on preterm birth.

Smoking and adverse socioeconomic characteristics are confirmed risk factors of preterm birth (Ahern et al. 2003; Ponce et al. 2005). Our findings of higher risk of preterm birth among mothers who were smoking and who lived at low SES areas provide further evidence for the role of smoking and socioeconomic inequity in the relationship between air pollution and preterm birth. It has been reported that multiple pregnancies and experience of previous pregnancy (especially for previous experience of preterm birth) are risk factors for preterm birth (Buchmayer et al. 2004; Kurdi et al. 2004). In this study, we propose that the experience of previous pregnancy and a multiple pregnancy may aggravate body's response to air pollution. Therefore, multiple pregnancies and those experiencing previous pregnancy should pay more attention to reduce the chance of exposure to air pollution to reduce the risk of preterm birth. So far, results from air pollution studies on the effect modification by gender are not uniform, and the biologic difference between boys and girls against air pollution remains unclear. A literature review summarised that male infants were generally less mature than females at term, and also at earlier gestational ages (Ghosh et al. 2007). This relatively less mature status may make them more vulnerable to risk factors like air pollution than females. However, gender analysis is essential in future research to elucidate this difference.

To our knowledge, this is the first study addressing the association between the very short-term (a few hours before onset of labour) air pollution exposure and the risk of preterm birth. The time-stratified case-crossover design was employed which ensured unbiased conditional logistic regression estimates and avoided time trend bias of the exposure (Janes et al. 2005). We ran one-pollutant and multiple-pollutant models to check the independent effect of pollutant, and we tailored the data by matching on specific groups to examine the effect modification of time-invariant factors. In addition, the threshold effect of air pollution was

indicated in our study which provided evidence of protecting people from bringing the pollutant concentration below the threshold point. A number of sensitivity analyses were performed to select the type and exposure time frame of hazardous pollutants.

Our study does have several limitations. First, we assigned mean pollutant concentrations of the specific hours' period preceding onset of labour to each preterm birth ignoring pregnant women's time spent indoors and outdoors. This may lead to exposure measurement error for the true exposure. However, we deducted the labour length when defined the time of labour onset to minimize this error. Moreover, this exposure measurement error can only underestimate the risk of preterm birth associated with increases in ambient air pollution concentrations (Zeger et al. 2000). Second, air pollution concentrations are likely to vary within and across urban study areas as a result of differences in meteorological, topographical, and environmental variables, and in the type and location of emission sources. We, as well as previous studies (Dominici et al. 2006; Guo et al. 2013; Peng et al. 2009; Samet et al. 2000), used city-wide average air pollutant concentrations to assign individual exposure which may bias effect estimates towards the null (Hutcheon et al. 2010). To improve the effect estimation, future studies should use air pollution exposure assessment tools with finer spatial resolution to characterize individual exposure such as land use regression models or interpolation methods. Third, as we didn't have individual-level SES data, SES linked to the living area during pregnancy was used to assess whether low SES level had higher risks of preterm birth associated with air pollution. This might underestimate the modification effect of SES. Fourth, our analyses were confined to a citywide dataset, resulting in the difficulty of generalizing the findings to other cities and other countries.

In conclusion, we found that sudden increases in the mean concentrations of ambient NO<sub>2</sub>, SO<sub>2</sub> and CO above the threshold levels in 24 or 48 hours immediately before onset of labour stimulated preterm birth. It provides latest evidence that reducing air pollution to a certain level or less could greatly benefit perinatal health, and that the influence may be quick effective. Some maternal demographic characteristics such as smoking and socioeconomic levels may modify the air pollution effects. We have proposed some biologic mechanisms underlying these associations.



## References

- Aagaard-Tillery KM, Nuthalapaty FS, Ramsey PS, Ramin KD. 2005. Preterm premature rupture of membranes: Perspectives surrounding controversies in management. *Am J Perinat* 22:287-297.
- Ahern J, Pickett KE, Selvin S, Abrams B. 2003. Preterm birth among African American and white women: A multilevel analysis of socioeconomic characteristics and cigarette smoking. *J Epidemiol Commun H* 57:606-611.
- Beck S, Wojdyla D, Say L, Betran AP, Merialdi M, Requejo JH, et al. 2010. The worldwide incidence of preterm birth: A systematic review of maternal mortality and morbidity. *B World Health Organ* 88:31-38.
- Blencowe H, Cousens S, Oestergaard MZ, Chou D, Moller A-B, Narwal R, et al. 2012. National, regional, and worldwide estimates of preterm birth rates in the year 2010 with time trends since 1990 for selected countries: A systematic analysis and implications. *Lancet* 379:2162-2172.
- Bobak M. 2000. Outdoor air pollution, low birth weight, and prematurity. *Environ Health Persp* 108:173.
- Buchmayer SM, Sparen P, Cnattingius S. 2004. Previous pregnancy loss: Risks related to severity of preterm delivery. *Am J Obstet Gynecol* 191:1225-1231.
- Chung JY, Honda Y, Hong YC, Pan XC, Guo Y, Kim H. 2009. Ambient temperature and mortality: An international study in four capital cities of East Asia. *Sci Total Environ* 408:390-396.
- Cunningham F, Leveno K, Bloom S, Spong CY, Dashe J. 2014. *Williams obstetrics* 24/e. E: McGraw Hill Professional.

- Dar-Row LA, Klein M, Correa A, Flanders WD, Waller L, Marcus M, et al. 2008. Ambient air pollution and preterm birth in atlanta, 1994-2004: A time-series analysis. *Birth Defects Res A* 82:367-367.
- Darrow LA, Klein M, Flanders WD, Waller LA, Correa A, Marcus M, et al. 2009. Ambient air pollution and preterm birth a time-series analysis. *Epidemiology* 20:689-698.
- Dominici F, Peng RD, Bell ML, Pham L, McDermott A, Zeger SL, et al. 2006. Fine particulate air pollution and hospital admission for cardiovascular and respiratory diseases. *JAMA* 295:1127-1134.
- Fleischer NL, Merialdi M, van Donkelaar A, Vadillo-Ortega F, Martin RV, Betran AP, et al. 2014. Outdoor air pollution, preterm birth, and low birth weight: Analysis of the World Health Organization Global Survey on Maternal and Perinatal Health. *Environ Health Persp* 122:425.
- Gehring U, Wijga AH, Fischer P, de Jongste JC, Kerkhof M, Koppelman GH, et al. 2011. Traffic-related air pollution, preterm birth and term birth weight in the piama birth cohort study. *Environ Res* 111:125-135.
- Ghosh R, Rankin J, Pless-Mulloli T, Glinianaia S. 2007. Does the effect of air pollution on pregnancy outcomes differ by gender? A systematic review. *Environ Res* 105:400-408.
- Goldenberg RL, Culhane JF, Iams JD, Romero R. 2008. Epidemiology and causes of preterm birth. *Lancet* 371:75-84.
- Guo Y, Barnett A, Yu W, Pan X, Ye X, Huang C, et al. 2011. A large change in temperature between neighbouring days increases the risk of mortality. *Plos One* 6:e16511.
- Guo Y, Li S, Tian Z, Pan X, Zhang J, Williams G. 2013. The burden of air pollution on years of life lost in Beijing, China, 2004-08: Retrospective regression analysis of daily deaths. *BMJ* 347:f7139.

- Hansen C, Neller A, Williams G, Simpson R. 2006. Maternal exposure to low levels of ambient air pollution and preterm birth in Brisbane, Australia. *BJOG: Int J Obstet Gy* 113:935-941.
- Hutcheon JA, Chiolero A, Hanley JA. 2010. Random measurement error and regression dilution bias. *BMJ* 340:c2289.
- Huynh M, Woodruff TJ, Parker JD, Schoendorf KC. 2006. Relationships between air pollution and preterm birth in California. *Paediatr Perinat Ep* 20:454-461.
- Jalaludin B, Mannes T, Morgan G, Lincoln D, Sheppard V, Corbett S. 2007. Impact of ambient air pollution on gestational age is modified by season in Sydney, Australia. *Environ Health* 6:16.
- Janes H, Sheppard L, Lumley T. 2005. Case–crossover analyses of air pollution exposure data: Referent selection strategies and their implications for bias. *Epidemiology* 16:717-726.
- Kan H, London SJ, Chen G, Zhang Y, Song G, Zhao N, et al. 2008. Season, sex, age, and education as modifiers of the effects of outdoor air pollution on daily mortality in Shanghai, China: The Public Health and Air Pollution in Asia (PAPA) Study. *Environ Health Perspect* 116:1183-1188.
- Kim H, Ha JS, Park J. 2006. High temperature, heat index, and mortality in 6 major cities in South Korea. *Arch Environ Occup Health* 61:265-270.
- Kurdi AM, Mesleh RA, Al-Hakeem MM, Khashoggi TY, Khalifa HM. 2004. Multiple pregnancy and preterm labor. *Saudi Med J* 25:632-637.
- Leem J-H, Kaplan BM, Shim YK, Pohl HR, Gotway CA, Bullard SM, et al. 2006. Exposures to air pollutants during pregnancy and preterm delivery. *Environ Health Persp* 114:905-910.

- Li S, Baker PJ, Jalaludin BB, Guo Y, Marks GB, Denison LS, et al. 2014. An Australian national panel study of diurnal temperature range and children's respiratory health. *Ann Allergy Asthma Immunol* 112:348-353 e341-348.
- Liu S, Krewski D, Shi Y, Chen Y, Burnett RT. 2003. Association between gaseous ambient air pollutants and adverse pregnancy outcomes in Vancouver, Canada. *Environ Health Persp* 111:1773.
- Llop S, Ballester F, Estarlich M, Esplugues A, Rebagliato M, Iñiguez C. 2010. Preterm birth and exposure to air pollutants during pregnancy. *Environ Res* 110:778-785.
- Lumley J. 2003. Defining the problem: The epidemiology of preterm birth. *BJOG: Int J Obstet Gy* 110:3-7.
- Lumley T, Levy D. 2000. Bias in the case-crossover design: Implications for studies of air pollution. *Environmetrics* 11:689-704.
- Olsson D, Ekstrom M, Forsberg B. 2012. Temporal variation in air pollution concentrations and preterm birth – A population based epidemiological study. *Int J Environ Res Public Health* 9:272-285.
- Peng RD, Bell ML, Geyh AS, McDermott A, Zeger SL, Samet JM, et al. 2009. Emergency admissions for cardiovascular and respiratory diseases and the chemical composition of fine particle air pollution. *Environ Health Perspect* 117:957-963.
- Ponce NA, Hoggatt KJ, Wilhelm M, Ritz B. 2005. Preterm birth: The interaction of traffic-related air pollution with economic hardship in Los Angeles neighborhoods. *Am J Epidemiol* 162:140-148.
- Qiu H, Yu IT-s, Tse LA, Tian L, Wang X, Wong TW. 2013. Is greater temperature change within a day associated with increased emergency hospital admissions for heart failure? *Circ Heart Fail* 6:930-935.

- R Core Team. 2015. A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
- Ritz B, Yu F, Chapa G, Fruin S. 2000. Effect of air pollution on preterm birth among children born in Southern California between 1989 and 1993. *Epidemiology* 11:502-511.
- Ritz B, Wilhelm M, Hoggatt KJ, Ghosh JKC. 2007. Ambient air pollution and preterm birth in the environment and pregnancy outcomes study at the University of California, Los Angeles. *Am J Epidemiol* 166:1045-1052.
- Sagiv SK, Mendola P, Loomis D, Herring AH, Neas LM, Savitz DA, et al. 2005. A time-series analysis of air pollution and preterm birth in Pennsylvania, 1997-2001. *Environ Health Perspect* 113:602-606.
- Samet JM, Dominici F, Currier FC, Coursac I, Zeger SL. 2000. Fine particulate air pollution and mortality in 20 US cities, 1987-1994. *New Engl J Med* 343:1742-1749.
- Slama R, Darrow L, Parker J, Woodruff T, Strickland M, Nieuwenhuijsen M, et al. 2008. Meeting report: Atmospheric pollution and human reproduction. *Environ Health Persp* 116:791-798.
- Suh Y-J, Ha E-H, Park H, Kim Y-J, Kim H, Hong Y-C. 2008. Gstm1 polymorphism along with PM10 exposure contributes to the risk of preterm delivery. *Mutat Res-Gen Tox En* 656:62-67.
- Suh YJ, Kim H, Seo JH, Park H, Kim YJ, Hong YC, et al. 2009. Different effects of PM10 exposure on preterm birth by gestational period estimated from time-dependent survival analyses. *Int Arch Occ Env Hea* 82:613-621.
- Wu J, Ren C, Delfino RJ, Chung J, Wilhelm M, Ritz B. 2009. Association between local traffic-generated air pollution and preeclampsia and preterm delivery in the South Coast air basin of California. *Environ Health Persp* 117:1773-1779.

- Yorifuji T, Naruse H, Kashima S, Murakoshi T, Doi H. 2015. Residential proximity to major roads and obstetrical complications. *Sci Total Environ* 508:188-192.
- Yu W, Vaneckova P, Mengersen K, Pan X, Tong S. 2010. Is the association between temperature and mortality modified by age, gender and socioeconomic status. *Sci Total Environ* 408:3513-3518.
- Zeger SL, Thomas D, Dominici F, Samet JM, Schwartz J, Dockery D, et al. 2000. Exposure measurement error in time-series studies of air pollution: Concepts and consequences. *Environ Health Persp* 108:419.
- Zeka A, Zanobetti A, Schwartz J. 2006. Individual-level modifiers of the effects of particulate matter on daily mortality. *Am J Epidemiol* 163:849-859.
- Zhao Q, Liang Z, Tao S, Zhu J, Du Y. 2011. Effects of air pollution on neonatal prematurity in Guangzhou of China: A time-series study. *Environ Health* 10:2.

Table 1: Characteristics of mothers and preterm births in Brisbane, Australia, 2009 – 2013.

Variables	N	%
Total	6949	100.0
Maternal age (years)		
< 35	5260	75.7
≥ 35	1689	24.3
Pre-pregnancy medical conditions		
No	5122	73.7
Yes	1827	26.3
Previous pregnancy		
0	2514	36.2
≥ 1	4435	63.8
Maternal smoking status		
Smokers	773	11.1
Non-smokers	6176	88.9
Number of births		
Single	4929	70.9
Multiple	2020	29.1
Baby gender		
Female	3276	47.1
Male	3673	52.9
Social-economic level		
Index 1 – 5 (Low development)	910	13.1
Index 6 – 10 (High development)	6039	86.9

Table 2: Summary statistics of hourly air pollution and weather conditions during 2009 – 2013 in Brisbane, Australia.

Variables	Mean $\pm$ SD	Percentiles				
		5th	25th	50th	75th	95th
PM2.5 ( $\mu\text{g}/\text{m}^3$ )	6.32 $\pm$ 4.05	3.20	4.35	5.55	7.24	11.10
PM10 ( $\mu\text{g}/\text{m}^3$ )	17.27 $\pm$ 15.33	9.78	13.08	15.79	18.94	26.29
NO2 (ppb)	6.52 $\pm$ 3.50	2.47	4.00	5.58	8.25	13.63
SO2 (ppb)	1.95 $\pm$ 3.26	0.42	0.75	1.15	2.00	5.36
O3 (ppb)	17.27 $\pm$ 8.59	5.33	10.56	16.11	23.00	32.89
CO (ppb)	219.30 $\pm$ 135.53	50.00	120.00	200.00	300.00	450.00
Temperature ( $^{\circ}\text{C}$ )	21.97 $\pm$ 4.83	13.25	18.78	22.48	25.46	29.24
Relative humidity (%)	70.82 $\pm$ 14.31	44.99	60.66	73.18	82.34	90.16

CO: carbon monoxide; NO2: nitrogen dioxide; O3: ozone; PM2.5: particulate matter 2.5 micrometres or less in diameter; PM10: particulate matter 10 micrometres or less in diameter; ppb: *parts per billion*; SD: standard division; SO2: sulphur dioxide;  $\mu\text{g}/\text{m}^3$ : microgram per cubic meter.



Table 3: The risks of preterm birth at 75th and 95th percentiles of air pollution against the minimum-preterm birth concentration of air pollution (threshold) in single-pollutant models.

	Threshold (ppb)	75th (ppb)	95th (ppb)	OR (95% CI)	
				75th VS Threshold	95th VS Threshold
NO <sub>2</sub>	7.6	8.3	11.3	1.01 (0.99, 1.03)	1.17 (1.08, 1.27)
SO <sub>2</sub>	3.8	2.2	4.8	1.04 (0.99, 1.08)	1.01 (0.99, 1.04)
CO	162.5	301.0	424.0	1.10 (1.01, 1.19)	1.18 (1.06, 1.32)

CI: confidence interval; CO: carbon monoxide (mean of 0 – 24 hours preceding onset of labour); NO<sub>2</sub>: nitrogen dioxide (mean of 0 – 24 hours preceding onset of labour); OR: odds ratio; ppb: *parts per billion*; SO<sub>2</sub>: sulphur dioxide (mean of 24 – 48 hours preceding onset of labour).

Table 4: The risks of preterm birth at 75th and 95th percentiles of air pollution against the minimum-preterm birth concentration of air pollution (threshold) in two-pollutant and three-pollutant models.

		OR (95% CI)	
		75th VS Threshold	95th VS Threshold
NO <sub>2</sub>	NO <sub>2</sub> + SO <sub>2</sub>	1.01 (0.99, 1.03)	1.17 (1.08, 1.27)
	NO <sub>2</sub> + CO	1.00 (0.98, 1.02)	1.13 (1.01, 1.26)
	NO <sub>2</sub> + SO <sub>2</sub> + CO	1.00 (0.98, 1.03)	1.13 (1.01, 1.26)
SO <sub>2</sub>	NO <sub>2</sub> + SO <sub>2</sub>	1.04 (0.99, 1.09)	1.01 (0.99, 1.04)
	SO <sub>2</sub> + CO	1.04 (0.99, 1.09)	1.01 (0.99, 1.04)
	NO <sub>2</sub> + SO <sub>2</sub> + CO	1.04 (0.99, 1.09)	1.01 (0.99, 1.04)
CO	NO <sub>2</sub> + CO	1.07 (0.97, 1.18)	1.11 (0.96, 1.27)
	SO <sub>2</sub> + CO	1.10 (1.01, 1.19)	1.18 (1.06, 1.32)
	NO <sub>2</sub> + SO <sub>2</sub> + CO	1.07 (0.97, 1.18)	1.11 (0.96, 1.27)

CI: confidence interval; CO: carbon monoxide (mean of 0 – 24 hours preceding onset of labour); NO<sub>2</sub>: nitrogen dioxide (mean of 0 – 24 hours preceding onset of labour); OR: odds ratio; ppb: *parts per billion*; SO<sub>2</sub>: sulphur dioxide (mean of 24 – 48 hours preceding onset of labour).

Table 5: Risks of preterm birth at 75th and 95th percentiles of air pollution against the minimum-preterm birth concentration of air pollution (threshold) by level of demographic factors. P-values for differences are presented in Supplemental Material, Table S3.

Factors		NO2 OR (95% CI)	SO2 OR (95% CI)	CO OR (95% CI)
Maternal age (years)				
< 35	75th VS Threshold	1.00 (0.99, 1.01)	1.06 (0.95, 1.17)	1.12 (0.94, 1.33)
	95th VS Threshold	1.26 (1.09, 1.47)	1.01 (0.97, 1.05)	1.30 (1.05, 1.61)
≥ 35	75th VS Threshold	1.01 (0.97, 1.06)	1.03 (0.98, 1.08)	1.09 (0.99, 1.20)
	95th VS Threshold	1.14 (1.03, 1.27)	1.02 (0.99, 1.05)	1.14 (1.01, 1.30)
Pre-pregnancy medical conditions				
No	75th VS Threshold	1.01 (0.99, 1.02)	1.04 (0.99, 1.10)	1.12 (1.02, 1.24)
	95th VS Threshold	1.18 (1.07, 1.30)	1.01 (0.99, 1.03)	1.24 (1.09, 1.41)
Yes	75th VS Threshold	1.67 (1.07, 2.61)	1.02 (0.95, 1.10)	1.11 (0.47, 2.59)
	95th VS Threshold	1.90 (1.18, 3.04)	1.03 (0.97, 1.09)	1.11 (0.51, 2.40)
Previous pregnancy				
0	75th VS Threshold	1.00 (0.99, 1.01)	1.10 (1.00, 1.22)	1.02 (0.93, 1.13)
	95th VS Threshold	1.06 (0.93, 1.21)	1.00 (0.99, 1.01)	1.09 (0.93, 1.26)
≥ 1	75th VS Threshold	1.01 (0.99, 1.04)	1.01 (0.97, 1.05)	1.15 (1.03, 1.29)
	95th VS Threshold	1.24 (1.12, 1.37)	1.03 (0.99, 1.08)	1.25 (1.09, 1.44)
Smoking				
Yes	75th VS Threshold	2.02 (1.05, 3.89)	1.08 (0.95, 1.24)	2.35 (0.58, 9.63)
	95th VS Threshold	2.59 (1.28, 5.25)	1.04 (0.97, 1.11)	2.47 (0.69, 8.82)
No	75th VS Threshold	1.01 (0.99, 1.03)	1.03 (0.98, 1.08)	1.06 (0.98, 1.15)
	95th VS Threshold	1.16 (1.06, 1.27)	1.01 (0.99, 1.04)	1.15 (1.03, 1.28)
Number of births				
Single	75th VS Threshold	1.00 (0.99, 1.01)	1.07 (0.99, 1.15)	1.09 (0.98, 1.21)
	95th VS Threshold	1.12 (1.02, 1.23)	1.00 (1.00, 1.00)	1.14 (1.00, 1.30)
Multiple	75th VS Threshold	1.03 (0.96, 1.10)	1.01 (0.97, 1.04)	1.11 (0.95, 1.28)
	95th VS Threshold	1.32 (1.12, 1.55)	1.10 (1.01, 1.19)	1.28 (1.06, 1.55)
Baby gender				
Female	75th VS Threshold	1.01 (0.98, 1.05)	1.01 (0.96, 1.05)	1.16 (1.00, 1.34)
	95th VS Threshold	1.20 (1.06, 1.36)	1.02 (0.97, 1.08)	1.26 (1.06, 1.49)
Male	75th VS Threshold	1.00 (0.99, 1.02)	1.07 (1.00, 1.14)	1.06 (0.96, 1.17)
	95th VS Threshold	1.15 (1.03, 1.28)	1.01 (0.99, 1.04)	1.14 (1.00, 1.30)
Social-economic level				
Index 1 – 5	75th VS Threshold	1.02 (0.99, 1.05)	1.05 (0.95, 1.17)	3.25 (1.72, 6.15)
	95th VS Threshold	1.19 (0.96, 1.46)	1.05 (0.98, 1.13)	3.27 (1.70, 6.27)
Index 6 – 10	75th VS Threshold	1.01 (0.99, 1.04)	1.03 (0.98, 1.09)	1.08 (0.99, 1.17)
	95th VS Threshold	1.18 (1.07, 1.29)	1.01 (0.99, 1.03)	1.17 (1.05, 1.30)

CI: confidence interval; CO: carbon monoxide (mean of 0 – 24 hours preceding onset of labour); NO2: nitrogen dioxide (mean of 0 – 24 hours preceding onset of labour); OR: odds ratio; SO2: sulphur dioxide (mean of 24 – 48 hours preceding onset of labour).

## Figure Legends

**Figure 1:** The relationships between air pollutants and preterm birth in single-pollutant models with 3 degrees of freedom natural cubic splines for air pollutants. CO: carbon monoxide; NO<sub>2</sub>: nitrogen dioxide; OR: odds ratio; ppb: *parts per billion*; SO<sub>2</sub>: sulphur dioxide.

**Figure 2:** The relationships between air pollutants and preterm birth in two-pollutant and three-pollutant models with 3 degrees of freedom natural cubic splines for air pollutants. (1) (4): NO<sub>2</sub> + SO<sub>2</sub>; (2) (7): NO<sub>2</sub> + CO; (5) (8): SO<sub>2</sub> + CO; (3) (6) (9): NO<sub>2</sub> + SO<sub>2</sub> + CO. CO: carbon monoxide; NO<sub>2</sub>: nitrogen dioxide; OR: odds ratio; ppb: *parts per billion*; SO<sub>2</sub>: sulphur dioxide.

Figure 1.




Figure 2.

